

Monoid hypersurfaces

Pål Hermunn Johansen, Magnus Løberg, Ragni Piene

Centre of Mathematics for Applications and Department of Mathematics
University of Oslo
P. O. Box 1053 Blindern
NO-0316 Oslo, Norway
{hermunn,mags,ragnip}@math.uio.no

Abstract. A monoid hypersurface is an irreducible hypersurface of degree d which has a singular point of multiplicity $d - 1$. Any monoid hypersurface admits a rational parameterization, hence is of potential interest in computer aided geometric design. We study properties of monoids in general and of monoid surfaces in particular. The main results include a description of the possible real forms of the singularities on a monoid surface other than the $(d - 1)$ -uple point. These results are applied to the classification of singularities on quartic monoid surfaces, complementing earlier work on the subject.

1 Introduction

A monoid hypersurface is an (affine or projective) irreducible algebraic hypersurface which has a singularity of multiplicity one less than the degree of the hypersurface. The presence of such a singular point forces the hypersurface to be rational: there is a rational parameterization given by (the inverse of) the linear projection of the hypersurface from the singular point.

The existence of an explicit rational parameterization makes such hypersurfaces potentially interesting objects in computer aided design. Moreover, since the “space” of monoids of a given degree is much smaller than the space of all hypersurfaces of that degree, one can hope to use monoids efficiently in (approximate or exact) implicitization problems. These were the reasons for considering monoids in the paper [17]. In [12] monoid curves are used to approximate other curves that are close to a monoid curve, and in [13] the same is done for monoid surfaces. In both articles the error of such approximations are analyzed – for each approximation, a bound on the distance from the monoid to the original curve or surface can be computed.

In this article we shall study properties of monoid hypersurfaces and the classification of monoid surfaces with respect to their singularities. Section 2 explores properties of monoid hypersurfaces in arbitrary dimension and over an arbitrary base field. Section 3 contains results on monoid surfaces, both over arbitrary fields and over $\mathbb{R}$. The last section deals with the classification of monoid surfaces of degree four. Real and complex quartic monoid surfaces were first studied by Rohn [15], who gave a fairly complete description of all possible cases. He also remarked [15, p. 56] that some of his results on quartic monoids hold for monoids of arbitrary degree; in particular, we believe he was aware of many of the results in Section 3. Takahashi, Watanabe, and

Higuchi [19] classify *complex* quartic monoid surfaces, but do not refer to Rohn. (They cite Jessop [7]; Jessop, however, only treats quartic surfaces with double points and refers to Rohn for the monoid case.) Here we aim at giving a short description of the possible singularities that can occur on quartic monoids, with special emphasis on the real case.

2 Basic properties

Let k be a field, let $\bar{k}$ denote its algebraic closure and $\mathbb{P}^n := \mathbb{P}_{\bar{k}}^n$ the projective n -space over $\bar{k}$. Furthermore we define the set of k -rational points $\mathbb{P}^n(k)$ as the set of points that admit representatives $(a_0 : \dots : a_n)$ with each $a_i \in k$.

For any homogeneous polynomial $F \in \bar{k}[x_0, \dots, x_n]$ of degree d and point $p = (p_0 : p_1 : \dots : p_n) \in \mathbb{P}^n$ we can define the multiplicity of $Z(F)$ at p . We know that $p_r \neq 0$ for some r , so we can assume $p_0 = 1$ and write

$$F = \sum_{i=0}^d x_0^{d-i} f_i(x_1 - p_1 x_0, x_2 - p_2 x_0, \dots, x_n - p_n x_0)$$

where f_i is homogeneous of degree i . Then the multiplicity of $Z(F)$ at p is defined to be the smallest i such that $f_i \neq 0$.

Let $F \in \bar{k}[x_0, \dots, x_n]$ be of degree $d \geq 3$. We say that the hypersurface $X = Z(F) \subset \mathbb{P}^n$ is a *monoid* hypersurface if X is irreducible and has a singular point of multiplicity $d - 1$.

In this article we shall only consider monoids $X = Z(F)$ where the singular point is k -rational. Modulo a projective transformation of $\mathbb{P}^n$ over k we may – and shall – therefore assume that the singular point is the point $O = (1 : 0 : \dots : 0)$.

Hence, we shall from now on assume that $X = Z(F)$, and

$$F = x_0 f_{d-1} + f_d,$$

where $f_i \in k[x_1, \dots, x_n] \subset k[x_0, \dots, x_n]$ is homogeneous of degree i and $f_{d-1} \neq 0$. Since F is irreducible, f_d is not identically 0, and f_{d-1} and f_d have no common (non-constant) factors.

The *natural rational parameterization* of the monoid $X = Z(F)$ is the map

$$\theta_F : \mathbb{P}^{n-1} \rightarrow \mathbb{P}^n$$

given by

$$\theta_F(a) = (f_d(a) : -f_{d-1}(a)a_1 : \dots : -f_{d-1}(a)a_n),$$

for $a = (a_1 : \dots : a_n)$ such that $f_{d-1}(a) \neq 0$ or $f_d(a) \neq 0$.

The set of lines through O form a $\mathbb{P}^{n-1}$. For every $a = (a_1 : \dots : a_n) \in \mathbb{P}^{n-1}$, the line

$$L_a := \{(s : ta_1 : \dots : ta_n) | (s : t) \in \mathbb{P}^1\} \quad (1)$$

intersects $X = Z(F)$ with multiplicity at least $d - 1$ in O . If $f_{d-1}(a) \neq 0$ or $f_d(a) \neq 0$, then the line L_a also intersects X in the point

$$\theta_F(a) = (f_d(a) : -f_{d-1}(a)a_1 : \dots : -f_{d-1}(a)a_n).$$

Hence the natural parameterization is the “inverse” of the projection of X from the point O . Note that θ_F maps $Z(f_{d-1}) \setminus Z(f_d)$ to O . The points where the parameterization map is not defined are called base points, and these points are precisely the common zeros of f_{d-1} and f_d . Each such point b corresponds to the line L_b contained in the monoid hypersurface. Additionally, every line of type L_b contained in the monoid hypersurface corresponds to a base point.

Note that $Z(f_{d-1}) \subset \mathbb{P}^{n-1}$ is the projective tangent cone to X at O , and that $Z(f_d)$ is the intersection of X with the hyperplane “at infinity” $Z(x_0)$.

Assume $P \in X$ is another singular point on the monoid X . Then the line L through P and O has intersection multiplicity at least $d - 1 + 2 = d + 1$ with X . Hence, according to Bezout’s theorem, L must be contained in X , so that this is only possible if $\dim X \geq 2$.

By taking the partial derivatives of F we can characterize the singular points of X in terms of f_d and f_{d-1} :

Lemma 1. *Let $\nabla = (\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n})$ be the gradient operator.*

- (i) *A point $P = (p_0 : p_1 : \dots : p_n) \in \mathbb{P}^n$ is singular on $Z(F)$ if and only if $f_{d-1}(p_1, \dots, p_n) = 0$ and $p_0 \nabla f_{d-1}(p_1, \dots, p_n) + \nabla f_d(p_1, \dots, p_n) = 0$.*
- (ii) *All singular points of $Z(F)$ are on lines L_a where a is a base point.*
- (iii) *Both $Z(f_{d-1})$ and $Z(f_d)$ are singular in a point $a \in \mathbb{P}^{n-1}$ if and only if all points on L_a are singular on X .*
- (iv) *If not all points on L_a are singular, then at most one point other than O on L_a is singular.*

Proof. (i) follows directly from taking the derivatives of $F = x_0 f_{d-1} + f_d$, and (ii) follows from (i) and the fact that $F(P) = 0$ for any singular point P . Furthermore, a point $(s : ta_1 : \dots : ta_n)$ on L_a is, by (i), singular if and only if

$$s \nabla f_{d-1}(ta) + \nabla f_d(ta) = t^{d-1}(s \nabla f_{d-1}(a) + t \nabla f_d(a)) = 0.$$

This holds for all $(s : t) \in \mathbb{P}^1$ if and only if $\nabla f_{d-1}(a) = \nabla f_d(a) = 0$. This proves (iii). If either $\nabla f_{d-1}(a)$ or $\nabla f_d(a)$ are nonzero, the equation above has at most one solution $(s_0 : t_0) \in \mathbb{P}^1$ in addition to $t = 0$, and (iv) follows.

Note that it is possible to construct monoids where $F \in k[x_0, \dots, x_n]$, but where no points of multiplicity $d - 1$ are k -rational. In that case there must be (at least) two such points, and the line connecting these will be of multiplicity $d - 2$. Furthermore, the natural parameterization will typically not induce a parameterization of the k -rational points from $\mathbb{P}^{n-1}(k)$.

3 Monoid surfaces

In the case of a monoid surface, the parameterization has a finite number of base points. From Lemma 1 (ii) we know that all singularities of the monoid other than O , are on lines L_a corresponding to these points. In what follows we will develop the theory for singularities on monoid surfaces — most of these results were probably known to Rohn [15, p. 56].

We start by giving a precise definition of what we shall mean by a monoid surface.

Definition 2. For an integer $d \geq 3$ and a field k of characteristic 0 the polynomials $f_{d-1} \in k[x_1, x_2, x_3]_{d-1}$ and $f_d \in k[x_1, x_2, x_3]_d$ define a normalized non-degenerate monoid surface $Z(F) \subset \mathbb{P}^3$, where $F = x_0 f_{d-1} + f_d \in k[x_0, x_1, x_2, x_3]$ if the following hold:

- (i) $f_{d-1}, f_d \neq 0$
- (ii) $\gcd(f_{d-1}, f_d) = 1$
- (iii) The curves $Z(f_{d-1}) \subset \mathbb{P}^2$ and $Z(f_d) \subset \mathbb{P}^2$ have no common singular point.

The curves $Z(f_{d-1}) \subset \mathbb{P}^2$ and $Z(f_d) \subset \mathbb{P}^2$ are called respectively the tangent cone and the intersection with infinity.

Unless otherwise stated, a surface that satisfies the conditions of Definition 2 shall be referred to simply as a *monoid surface*.

Since we have finitely many base points b and each line L_b contains at most one singular point in addition to O , monoid surfaces will have only finitely many singularities, so all singularities will be isolated. (Note that Rohn included surfaces with nonisolated singularities in his study [15].) We will show that the singularities other than O can be classified by local intersection numbers.

Definition 3. Let $f, g \in k[x_1, x_2, x_3]$ be nonzero and homogeneous. Assume $p = (p_1 : p_2 : p_3) \in Z(f, g) \subset \mathbb{P}^2$, and define the local intersection number

$$I_p(f, g) = \lg \frac{\bar{k}[x_1, x_2, x_3]_{m_p}}{(f, g)},$$

where $\bar{k}$ is the algebraic closure of k , $m_p = (p_2x_1 - p_1x_2, p_3x_1 - p_1x_3, p_3x_2 - p_2x_3)$ is the homogeneous ideal of p , and $\lg$ denotes the length of the local ring as a module over itself.

Note that $I_p(f, g) \geq 1$ if and only if $f(p) = g(p) = 0$. When $I_p(f, g) = 1$ we say that f and g intersect transversally at p . The terminology is justified by the following lemma:

Lemma 4. Let $f, g \in k[x_1, x_2, x_3]$ be nonzero and homogeneous and $p \in Z(f, g)$. Then the following are equivalent:

- (i) $I_p(f, g) > 1$
- (ii) f is singular at p , g is singular at p , or $\nabla f(p)$ and $\nabla g(p)$ are nonzero and parallel.
- (iii) $s\nabla f(p) + t\nabla g(p) = 0$ for some $(s, t) \neq (0, 0)$

Proof. (ii) is equivalent to (iii) by a simple case study: f is singular at p if and only if (iii) holds for $(s, t) = (1, 0)$, g is singular at p if and only if (iii) holds for $(s, t) = (0, 1)$, and $\nabla f(p)$ and $\nabla g(p)$ are nonzero and parallel if and only if (iii) holds for some $s, t \neq 0$.

We can assume that $p = (0 : 0 : 1)$, so $I_p(f, g) = \lg S$ where

$$S = \frac{\bar{k}[x_1, x_2, x_3]_{(x_1, x_2)}}{(f, g)}.$$

Furthermore, let $d = \deg f$, $e = \deg g$ and write

$$f = \sum_{i=1}^d f_i x_3^{d-i} \text{ and } g = \sum_{i=1}^e g_i x_3^{e-i}$$

where f_i, g_i are homogeneous of degree i .

If f is singular at p , then $f_1 = 0$. Choose $\ell = ax_1 + bx_2$ such that ℓ is not a multiple of g_1 . Then ℓ will be a nonzero non-invertible element of S , so the length of S is greater than 1.

We have $\nabla f(p) = (\nabla f_1(p), 0)$ and $\nabla g(p) = (\nabla g_1(p), 0)$. If they are parallel, choose $\ell = ax_0 + bx_1$ such that ℓ is not a multiple of f_1 (or g_1), and argue as above.

Finally assume that f and g intersect transversally at p . We may assume that $f_1 = x_1$ and $g_1 = x_2$. Then $(f, g) = (x_1, x_2)$ as ideals in the local ring $\bar{k}[x_1, x_2, x_3]_{(x_1, x_2)}$. This means that S is isomorphic to the field $\bar{k}(x_3)$. The length of any field is 1, so $I_p(f, g) = \lg S = 1$.

Now we can say which are the lines L_b , with $b \in Z(f_{d-1}, f_d)$, that contain a singularity other than O :

Lemma 5. *Let f_{d-1} and f_d be as in Definition 2. The line L_b contains a singular point other than O if and only if $Z(f_{d-1})$ is nonsingular at b and the intersection multiplicity $I_b(f_{d-1}, f_d) > 1$.*

Proof. Let $b = (b_1 : b_2 : b_3)$ and assume that $(b_0 : b_1 : b_2 : b_3)$ is a singular point of $Z(F)$. Then, by Lemma 1, $f_{d-1}(b_1, b_2, b_3) = f_d(b_1, b_2, b_3) = 0$ and $b_0 \nabla f_{d-1}(b_1, b_2, b_3) + \nabla f_d(b_1, b_2, b_3) = 0$, which implies $I_b(f_{d-1}, f_d) > 1$. Furthermore, if f_{d-1} is singular at b , then the gradient $\nabla f_{d-1}(b_1, b_2, b_3) = 0$, so f_d , too, is singular at b , contrary to our assumptions.

Now assume that $Z(f_{d-1})$ is nonsingular at $b = (b_1 : b_2 : b_3)$ and the intersection multiplicity $I_b(f_{d-1}, f_d) > 1$. The second assumption implies $f_{d-1}(b_1, b_2, b_3) = f_d(b_1, b_2, b_3) = 0$ and $s \nabla f_{d-1}(b_1, b_2, b_3) = t \nabla f_d(b_1, b_2, b_3)$ for some $(s, t) \neq (0, 0)$. Since $Z(f_{d-1})$ is nonsingular at b , we know that $\nabla f_{d-1}(b_1, b_2, b_3) \neq 0$, so $t \neq 0$. Now $(-s/t : b_1 : b_2 : b_3) \neq (1 : 0 : 0 : 0)$ is a singular point of $Z(F)$ on the line L_b .

Recall that an A_n singularity is a singularity with normal form $x_1^2 + x_2^2 + x_3^{n+1}$, see [3, p. 184].

Proposition 6. *Let f_{d-1} and f_d be as in Definition 2, and assume $P = (p_0 : p_1 : p_2 : p_3) \neq (1 : 0 : 0 : 0)$ is a singular point of $Z(F)$ with $I_{(p_1:p_2:p_3)}(f_{d-1}, f_d) = m$. Then P is an A_{m-1} singularity.*

Proof. We may assume that $P = (0 : 0 : 0 : 1)$ and write the local equation

$$g := F(x_0, x_1, x_2, 1) = x_0 f_{d-1}(x_1, x_2, 1) + f_d(x_1, x_2, 1) = \sum_{i=2}^d g_i \quad (2)$$

with $g_i \in \bar{k}[x_0, x_1, x_2]$ homogeneous of degree i . Since $Z(f_{d-1})$ is nonsingular at $O := (0 : 0 : 1)$, we can assume that the linear term of $f_{d-1}(x_1, x_2, 1)$ is equal to x_1 .

The quadratic term g_2 of g is then $g_2 = x_0x_1 + ax_1^2 + bx_1x_2 + cx_2^2$ for some $a, b, c \in k$. The Hessian matrix of g evaluated at P is

$$H(g)(0, 0, 0) = H(g_2)(0, 0, 0) = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 2a & b \\ 0 & b & 2c \end{pmatrix}$$

which has corank 0 when $c \neq 0$ and corank 1 when $c = 0$. By [3, p. 188], P is an A_1 singularity when $c \neq 0$ and an A_n singularity for some n when $c = 0$.

The index n of the singularity is equal to the Milnor number

$$\mu = \dim_{\bar{k}} \frac{\bar{k}[x_0, x_1, x_2]_{(x_0, x_1, x_2)}}{J_g} = \dim_{\bar{k}} \frac{\bar{k}[x_0, x_1, x_2]_{(x_0, x_1, x_2)}}{\left(\frac{\partial g}{\partial x_0}, \frac{\partial g}{\partial x_1}, \frac{\partial g}{\partial x_2} \right)}.$$

We need to show that $\mu = I_0(f_{d-1}, f_d) - 1$. From the definition of the intersection multiplicity, it is not hard to see that

$$I_0(f_{d-1}, f_d) = \dim_{\bar{k}} \frac{\bar{k}[x_1, x_2]_{(x_1, x_2)}}{(f_{d-1}(x_1, x_2, 1), f_d(x_1, x_2, 1))}.$$

The singularity at p is isolated, so the Milnor number is finite. Furthermore, since $\gcd(f_{d-1}, f_d) = 1$, the intersection multiplicity is finite. Therefore both dimensions can be calculated in the completion rings. For the rest of the proof we view f_{d-1} and f_d as elements of the power series rings $\bar{k}[[x_1, x_2]] \subset \bar{k}[[x_0, x_1, x_2]]$, and all calculations are done in these rings.

Since $Z(f_{d-1})$ is smooth at O , we can write

$$f_{d-1}(x_1, x_2, 1) = (x_1 - \phi(x_2)) u(x_1, x_2)$$

for some power series $\phi(x_2)$ and invertible power series $u(x_1, x_2)$. To simplify notation we write $u = u(x_1, x_2) \in \bar{k}[[x_1, x_2]]$.

The Jacobian ideal J_g is generated by the three partial derivatives:

$$\begin{aligned} \frac{\partial g}{\partial x_0} &= (x_1 - \phi(x_2)) u \\ \frac{\partial g}{\partial x_1} &= x_0 \left(u + (x_1 - \phi(x_2)) \frac{\partial u}{\partial x_1} \right) + \frac{\partial f_d}{\partial x_1}(x_1, x_2) \\ \frac{\partial g}{\partial x_2} &= x_0 \left(-\phi'(x_2)u + (x_1 - \phi(x_2)) \frac{\partial u}{\partial x_2} \right) + \frac{\partial f_d}{\partial x_2}(x_1, x_2) \end{aligned}$$

By using the fact that $x_1 - \phi(x_2) \in \left(\frac{\partial g}{\partial x_0} \right)$ we can write J_g without the symbols $\frac{\partial u}{\partial x_1}$ and $\frac{\partial u}{\partial x_2}$:

$$J_g = \left(x_1 - \phi(x_2), x_0u + \frac{\partial f_d}{\partial x_1}(x_1, x_2), -x_0u\phi'(x_2) + \frac{\partial f_d}{\partial x_2}(x_1, x_2) \right)$$

To make the following calculations clear, define the polynomials h_i by writing $f_d(x_1, x_2, 1) = \sum_{i=0}^d x_1^i h_i(x_2)$. Now

$$J_g = \left(x_1 - \phi(x_2), x_0 u + \sum_{i=1}^d i x_1^{i-1} h_i(x_2), -x_0 u \phi'(x_2) + \sum_{i=0}^d x_1^i h'_i(x_2) \right),$$

so

$$\frac{\bar{k}[[x_0, x_1, x_2]]}{J_g} = \frac{\bar{k}[[x_2]]}{(A(x_2))}$$

where

$$A(x_2) = \phi'(x_2) \left(\sum_{i=1}^d i \phi(x_2)^{i-1} h_i(x_2) \right) + \left(\sum_{i=0}^d \phi(x_2)^i h'_i(x_2) \right).$$

For the intersection multiplicity we have

$$\frac{\bar{k}[[x_1, x_2]]}{\left(f_{d-1}(x_1, x_2, 1), f_d(x_1, x_2, 1) \right)} = \frac{\bar{k}[[x_1, x_2]]}{\left(x_1 - \phi(x_2), \sum_{i=0}^d x_1^i h_i(x_2) \right)} = \frac{\bar{k}[[x_2]]}{(B(x_2))}$$

where $B(x_2) = \sum_{i=0}^d \phi(x_2)^i h_i(x_2)$. Observing that $B'(x_2) = A(x_2)$ gives the result $\mu = I_0(f_{d-1}, f_d) - 1$.

Corollary 7. *A monoid surface of degree d can have at most $\frac{1}{2}d(d-1)$ singularities in addition to O . If this number of singularities is obtained, then all of them will be of type A_1 .*

Proof. The sum of all local intersection numbers $I_a(f_{d-1}, f_d)$ is given by Bézout's theorem:

$$\sum_{a \in \mathbb{Z}(f_{d-1}, f_d)} I_a(f_{d-1}, f_d) = d(d-1).$$

The line L_a will contain a singularity other than O only if $I_a(f_{d-1}, f_d) \geq 2$, giving a maximum of $\frac{1}{2}d(d-1)$ singularities in addition to O . Also, if this number is obtained, all local intersection numbers must be exactly 2, so all singularities other than O will be of type A_1 .

Both Proposition 6 and Corollary 7 were known to Rohn, who stated these results only in the case $d = 4$, but said they could be generalized to arbitrary d [15, p. 60].

For the rest of the section we will assume $k = \mathbb{R}$. It turns out that we can find a *real* normal form for the singularities other than O . The complex singularities of type A_n come in several real types, with normal forms $x_1^2 \pm x_2^2 \pm x_3^{n+1}$. Varying the $\pm$ gives two types for $n = 1$ and n even, and three types for $n \geq 3$ odd. The real type with normal form $x_1^2 - x_2^2 + x_3^{n+1}$ is called an A_n^- singularity, or of type A^- , and is what we find on real monoids:

Proposition 8. *On a real monoid, all singularities other than O are of type A^- .*

Proof. Assume $p = (0 : 0 : 1)$ is a singular point on $Z(F)$ and set $g = F(x_0, x_1, x_2, 1)$ as in the proof of Proposition 6.

First note that $u^{-1}g = x_0(x_1 - \phi(x_2)) + f_d(x_1, x_2)u^{-1}$ is an equation for the singularity. We will now prove that $u^{-1}g$ is right equivalent to $\pm(x_0^2 - x_1^2 + x_2^n)$, for some n , by constructing right equivalent functions $u^{-1}g =: g_{(0)} \sim g_{(1)} \sim g_{(2)} \sim g_{(3)} \sim \pm(x_0^2 - x_1^2 + x_2^n)$. Let

$$\begin{aligned} g_{(1)}(x_0, x_1, x_2) &= g_{(0)}(x_0, x_1 + \phi(x_2), x_2) \\ &= x_0x_1 + f_d(x_1 + \phi(x_2), x_2)u^{-1}(x_1 + \phi(x_2), x_2) \\ &= x_0x_1 + \psi(x_1, x_2) \end{aligned}$$

where $\psi(x_1, x_2) \in \mathbb{R}[[x_1, x_2]]$. Write $\psi(x_1, x_2) = x_1\psi_1(x_1, x_2) + \psi_2(x_2)$ and define

$$g_{(2)}(x_0, x_1, x_2) = g_{(1)}(x_0 - \psi_1(x_1, x_2), x_1, x_2) = x_0x_1 + \psi_2(x_2).$$

The power series $\psi_2(x_2)$ can be written on the form

$$\psi_2(x_2) = sx_2^n(a_0 + a_1x_2 + a_2x_2^2 + \dots)$$

where $s = \pm 1$ and $a_0 > 0$. We see that $g_{(2)}$ is right equivalent to $g_{(3)} = x_0x_1 + sx_2^n$ since

$$g_{(2)}(x_0, x_1, x_2) = g_{(3)}\left(x_0, x_1, x_2 \sqrt[n]{a_0 + a_1x_2 + a_2x_2^2 + \dots}\right).$$

Finally we see that

$$g_{(4)}(x_0, x_1, x_2) := g_{(3)}(sx_0 - sx_1, x_0 + x_1, x_2) = s(x_0^2 - x_1^2 + x_2^n)$$

proves that $u^{-1}g$ is right equivalent to $s(x_0^2 - x_1^2 + x_2^n)$ which is an equation for an A_{n-1} singularity with normal form $x_0^2 - x_1^2 + x_2^n$.

Note that for $d = 3$, the singularity at O can be an A_1^+ singularity. This happens for example when $f_2 = x_0^2 + x_1^2 + x_2^2$.

For a *real* monoid, Corollary 7 implies that we can have at most $\frac{1}{2}d(d-1)$ *real* singularities in addition to O . We can show that the bound is sharp by a simple construction:

Example. To construct a monoid with the maximal number of real singularities, it is sufficient to construct two *affine* real curves in the xy -plane defined by equations f_{d-1} and f_d of degrees $d-1$ and d such that the curves intersect in $d(d-1)/2$ points with multiplicity 2. Let $m \in \{d-1, d\}$ be odd and set

$$f_m = \varepsilon - \prod_{i=1}^m \left(x \sin\left(\frac{2i\pi}{m}\right) + y \cos\left(\frac{2i\pi}{m}\right) + 1 \right).$$

For $\varepsilon > 0$ sufficiently small there exist at least $\frac{m+1}{2}$ radii $r > 0$, one for each root of the univariate polynomial $f_m|_{x=0}$, such that the circle $x^2 + y^2 - r^2$ intersects f_m in m points




Fig. 1. The curves f_m for $m = 3, 5$ and corresponding circles

with multiplicity 2. Let f_{2d-1-m} be a product of such circles. Now the homogenizations of f_{d-1} and f_d define a monoid surface with $1 + \frac{1}{2}d(d-1)$ singularities. See Figure 1.

Proposition 6 and Bezout's theorem imply that the maximal Milnor number of a singularity other than O is $d(d-1) - 1$. The following example shows that this bound can be achieved on a real monoid:

Example. The surface $X \subset \mathbb{P}^3$ defined by $F = x_0(x_1x_2^{d-2} + x_3^{d-1}) + x_1^d$ has exactly two singular points. The point $(1 : 0 : 0 : 0)$ is a singularity of multiplicity $d-1$ with Milnor number $\mu = (d^2 - 3d + 1)(d-2)$, while the point $(0 : 0 : 1 : 0)$ is an $A_{d(d-1)-1}$ singularity. A picture of this surface for $d = 4$ is shown in Figure 2.

4 Quartic monoid surfaces

Every cubic surface with isolated singularities is a monoid. Both smooth and singular cubic surfaces have been studied extensively, most notably in [16], where real cubic surfaces and their singularities were classified, and more recently in [18], [4], and [8]. The site [9] contains additional pictures and references.

In this section we shall consider the case $d = 4$. The classification of real and complex *quartic* monoid surfaces was started by Rohn [15]. (In addition to considering the singularities, Rohn studied the existence of lines *not* passing through the triple point, and that of other special curves on the monoid.) In [19], Takahashi, Watanabe, and Higuchi described the singularities of such *complex* surfaces. The monoid singularity of a quartic monoid is minimally elliptic [21], and minimally elliptic singularities have the same complex topological type if and only if their dual graphs are isomorphic [10]. In [10] all possible dual graphs for minimally elliptic singularities are listed, along with example equations.

Using Arnold's notation for the singularities, we use and extend the approach of Takahashi, Watanabe, and Higuchi in [19].




Fig. 2. The surface defined by $F = x_0(x_1x_2^{d-2} + x_3^{d-1}) + x_1^d$ for $d = 4$.

Consider a quartic monoid surface, $X = Z(F)$, with $F = x_0f_3 + f_4$. The tangent cone, $Z(f_3)$, can be of one of nine (complex) types, each needing a separate analysis.

For each type we fix f_3 , but any other tangent cone of the same type will be projectively equivalent (over the complex numbers) to this fixed f_3 . The nine different types are:

1. Nodal irreducible curve, $f_3 = x_1x_2x_3 + x_2^3 + x_3^3$.
2. Cuspidal curve, $f_3 = x_1^3 - x_2^2x_3$.
3. Conic and a chord, $f_3 = x_3(x_1x_2 + x_3^2)$.
4. Conic and a tangent line, $f_3 = x_3(x_1x_3 + x_2^2)$.
5. Three general lines, $f_3 = x_1x_2x_3$.
6. Three lines meeting in a point, $f_3 = x_2^3 - x_2x_3^2$.
7. A double line and another line, $f_3 = x_2x_3^2$.
8. A triple line $f_3 = x_3^3$.
9. A smooth curve, $f_3 = x_1^3 + x_2^3 + x_3^3 + 3ax_0x_1x_3$ where $a^3 \neq -1$.

To each quartic monoid we can associate, in addition to the type, several integer invariants, all given as intersection numbers. From [19] we know that, for the types 1–3, 5, and 9, these invariants will determine the singularity type of O up to right equivalence. In the other cases the singularity series, as defined by Arnol'd in [1] and [2], is determined by the type of f_3 . We shall use, without proof, the results on the singularity type of O due to [19]; however, we shall use the notations of [1] and [2].

We complete the classification begun in [19] by supplying a complete list of the possible singularities occurring on a quartic monoid. In addition, we extend the results to the case of *real* monoids. Our results are summarized in the following theorem.

Theorem 9. *On a quartic monoid surface, singularities other than the monoid point can occur as given in Table 1. Moreover, all possibilities are realizable on real quartic*

monoids with a real monoid point, and with the other singularities being real and of type A^- .

Case	Triple point	Invariants and constraints	Other singularities
1	$T_{3,3,4}$		$A_{m_i-1}, \sum m_i = 12$
	$T_{3,3,3+m}$	$m = 2, \dots, 12$	$A_{m_i-1}, \sum m_i = 12 - m$
2	Q_{10}		$A_{m_i-1}, \sum m_i = 12$
	T_{9+m}	$m = 2, 3$	$A_{m_i-1}, \sum m_i = 12 - m$
3	$T_{3,4+r_0,4+r_1}$	$r_0 = \max(j_0, k_0), r_1 = \max(j_1, k_1),$ $j_0 > 0 \leftrightarrow k_0 > 0, \min(j_0, k_0) \leq 1,$ $j_1 > 0 \leftrightarrow k_1 > 0, \min(j_1, k_1) \leq 1$	$A_{m_i-1}, \sum m_i = 4 - k_0 - k_1,$ $A_{m'_i-1}, \sum m'_i = 8 - j_0 - j_1$
4	S series	$j_0 \leq 8, k_0 \leq 4, \min(j_0, k_0) \leq 2,$ $j_0 > 0 \leftrightarrow k_0 > 0, j_1 > 0 \leftrightarrow k_0 > 1$	$A_{m_i-1}, \sum m_i = 4 - k_0,$ $A_{m'_i-1}, \sum m'_i = 8 - j_0$
5	$T_{4+j_k,4+j_l,4+j_m}$	$m_1 + l_1 \leq 4, k_2 + m_2 \leq 4,$ $k_3 + l_3 \leq 4, k_2 > 0 \leftrightarrow k_3 > 0,$ $l_1 > 0 \leftrightarrow l_3 > 0, m_1 > 0 \leftrightarrow m_2 > 0,$ $\min(k_2, k_3) \leq 1, \min(l_1, l_3) \leq 1,$ $\min(m_1, m_2) \leq 1, j_k = \max(k_2, k_3),$ $j_l = \max(l_1, l_3), j_m = \max(m_1, m_2)$	$A_{m_i-1}, \sum m_i = 4 - m_1 - l_1,$ $A_{m'_i-1}, \sum m'_i = 4 - k_2 - m_2,$ $A_{m''_i-1}, \sum m''_i = 4 - k_3 - l_3$
6	U series	$j_1 > 0 \leftrightarrow j_2 > 0 \leftrightarrow j_3 > 0,$ at most one of $j_1, j_2, j_3 > 1,$ $j_1, j_2, j_3 \leq 4$	$A_{m_i-1}, \sum m_i = 4 - j_1,$ $A_{m'_i-1}, \sum m'_i = 4 - j_2,$ $A_{m''_i-1}, \sum m''_i = 4 - j_3$
7	V series	$j_0 > 0 \leftrightarrow k_0 > 0, \min(j_0, k_0) \leq 1,$ $j_0 \leq 4, k_0 \leq 4$	$A_{m_i-1}, \sum m_i = 4 - j_0,$
8	V' series		None
9	$P_8 = T_{3,3,3}$		$A_{m_i-1}, \sum m_i = 12$

Table 1. Possible configurations of singularities for each case

Proof. The invariants listed in the “Invariants and constraints” column are all nonnegative integers, and any set of integer values satisfying the equations represents one possible set of invariants, as described above. Then, for each set of invariants, (positive) intersection multiplicities, denoted m_i , m'_i and m''_i , will determine the singularities other than O . The column “Other singularities” give these and the equations they must satisfy. Here we use the notation A_0 for a line L_a on $Z(F)$ where O is the only singular point.

The analyses of the nine cases share many similarities, and we have chosen not to go into great detail when one aspect of a case differs little from the previous one. We end the section with a discussion on the possible real forms of the tangent cone and how this affects the classification of the *real* quartic monoids.

In all cases, we shall write

$$\begin{aligned} f_4 = & a_1x_1^4 + a_2x_1^3x_2 + a_3x_1^3x_3 + a_4x_1^2x_2^2 + a_5x_1^2x_2x_3 \\ & + a_6x_1^2x_3^2 + a_7x_1x_2^3 + a_8x_1x_2^2x_3 + a_9x_1x_2x_3^2 + a_{10}x_1x_3^3 \\ & + a_{11}x_2^4 + a_{12}x_2^3x_3 + a_{13}x_2^2x_3^2 + a_{14}x_2x_3^3 + a_{15}x_3^4 \end{aligned}$$

and we shall investigate how the coefficients $a_1, \dots, a_{15}$ are related to the geometry of the monoid.

Case 1. The tangent cone is a nodal irreducible curve, and we can assume

$$f_3(x_1, x_2, x_3) = x_1x_2x_3 + x_2^3 + x_3^3.$$

The nodal curve is singular at $(1 : 0 : 0)$. If $f_4(1, 0, 0) \neq 0$, then O is a $T_{3,3,4}$ singularity [19]. We recall that $(1 : 0 : 0)$ cannot be a singular point on $Z(f_4)$ as this would imply a singular line on the monoid, so we assume that either $(1 : 0 : 0) \notin Z(f_4)$ or $(1 : 0 : 0)$ is a smooth point on $Z(f_4)$. Let m denote the intersection number $I_{(1:0:0)}(f_3, f_4)$. Since $Z(f_3)$ is singular at $(1 : 0 : 0)$ we have $m \neq 1$. From [19] we know that O is a $T_{3,3,3+m}$ singularity for $m = 2, \dots, 12$. Note that some of these complex singularities have two real forms, as illustrated in Figure 3.




Fig. 3. The monoids $Z(x^3 + y^3 + 5xyz - z^3(x + y))$ and $Z(x^3 + y^3 + 5xyz - z^3(x - y))$ both have a $T_{3,3,5}$ singularity, but the singularities are not right equivalent over $\mathbb{R}$. (The pictures are generated by the program [5].)

Bézout's theorem and Proposition 6 limit the possible configurations of singularities on the monoid for each m . Let $\theta(s, t) = (-s^3 - t^3, s^2t, st^2)$. Then the tangent cone $Z(f_3)$ is parameterized by θ as a map from $\mathbb{P}^1$ to $\mathbb{P}^2$. When we need to compute the intersection numbers between the rational curve $Z(f_3)$ and the curve $Z(f_4)$, we can do

that by studying the roots of the polynomial $f_4(\theta)$. Expanding the polynomial gives

$$\begin{aligned} f_4(\theta)(s, t) = & a_1 s^{12} - a_2 s^{11} t + (-a_3 + a_4) s^{10} t^2 + (4a_1 + a_5 - a_7) s^9 t^3 \\ & + (-3a_2 + a_6 - a_8 + a_{11}) s^8 t^4 + (-3a_3 + 2a_4 - a_9 + a_{12}) s^7 t^5 \\ & + (6a_1 + 2a_5 - a_7 - a_{10} + a_{13}) s^6 t^6 \\ & + (-3a_2 + 2a_6 - a_8 + a_{14}) s^5 t^7 + (-3a_3 + a_4 - a_9 + a_{15}) s^4 t^8 \\ & + (4a_1 + a_5 - a_{10}) s^3 t^9 + (-a_2 + a_6) s^2 t^{10} - a_3 s t^{11} + a_1 t^{12}. \end{aligned}$$

This polynomial will have roots at $(0 : 1)$ and $(1 : 0)$ if and only if $f_4(1, 0, 0) = a_1 = 0$. When $a_1 = 0$ we may (by symmetry) assume $a_3 \neq 0$, so that $(0 : 1)$ is a simple root and $(1 : 0)$ is a root of multiplicity $m - 1$. Other roots of $f_4(\theta)$ correspond to intersections of $Z(f_3)$ and $Z(f_4)$ away from $(1 : 0 : 0)$. The multiplicity m_i of each root is equal to the corresponding intersection multiplicity, giving rise to an A_{m_i-1} singularity if $m_i > 0$, as described by Proposition 6, or a line $L_a \subset Z(F)$ with O as the only singular point if $m_i = 1$.

The polynomial $f_4(\theta)$ defines a linear map from the coefficient space k^{15} of f_4 to the space of homogeneous polynomials of degree 12 in s and t . By elementary linear algebra, we see that the image of this map is the set of polynomials of the form

$$b_0 s^{12} + b_1 s^{11} t + b_2 s^{10} t^2 + \cdots + b_{12} t^{12}$$

where $b_0 = b_{12}$. The kernel of the map corresponds to the set of polynomials of the form ℓf_3 where ℓ is a linear form. This means that $f_4(\theta) \equiv 0$ if and only if f_3 is a factor in f_4 , making $Z(F)$ reducible and not a monoid.

For every $m = 0, 2, 3, 4, \dots, 12$ we can select r parameter points

$$p_1, \dots, p_r \in \mathbb{P}^1 \setminus \{(0 : 1), (1 : 0)\}$$

and positive multiplicities $m_1, \dots, m_r$ with $m_1 + \cdots + m_r = 12 - m$ and try to describe the polynomials f_4 such that $f_4(\theta)$ has a root of multiplicity m_i at p_i for each $i = 1, \dots, r$.

Still assuming $a_3 \neq 0$ whenever $a_1 = 0$, any such choice of parameter points $p_1, \dots, p_r$ and multiplicities $m_1, \dots, m_r$ corresponds to a polynomial $q = b_0 s^{12} + b_1 s^{11} t + \cdots + b_{12} t^{12}$ that is, up to a nonzero constant, uniquely determined.

Now, q is equal to $f_4(\theta)$ for some f_4 if and only if $b_0 = b_{12}$. If $m \geq 2$, then q contains a factor st^{m-1} , so $b_0 = b_{12} = 0$, giving $q = f_4(\theta)$ for some f_4 . In fact, when $m \geq 2$ any choice of $p_1, \dots, p_r$ and $m_1, \dots, m_r$ with $m_1 + \cdots + m_r = 12 - m$ corresponds to a four dimensional space of equations f_4 that gives this set of roots and multiplicities in $f_4(\theta)$. If f'_4 is one such f_4 , then any other is of the form $\lambda f'_4 + \ell f_3$ for some constant $\lambda \neq 0$ and linear form ℓ . All of these give monoids that are projectively equivalent.

When $m = 0$, we write $p_i = (\alpha_i : \beta_i)$ for $i = 1, \dots, r$. The condition $b_0 = b_{12}$ on the coefficients of q translates to

$$\alpha_1^{m_1} \cdots \alpha_r^{m_r} = \beta_1^{m_1} \cdots \beta_r^{m_r}. \quad (3)$$

This means that any choice of parameter points $(\alpha_1 : \beta_1), \dots, (\alpha_r : \beta_r)$ and multiplicities $m_1, \dots, m_r$ with $m_1 + \dots + m_r = 12$ that satisfy condition (3) corresponds to a four dimensional family $\lambda f'_4 + \ell f_3$, giving a unique monoid up to projective equivalence.

For example, we can have an A_{11} singularity only if $f_4(\theta)$ is of the form $(\alpha s - \beta t)^{12}$. Condition (3) implies that this can only happen for 12 parameter points, all of the form $(1 : \omega)$, where $\omega^{12} = 1$. Each such parameter point $(1 : \omega)$ corresponds to a monoid uniquely determined up to projective equivalence. However, since there are six projective transformations of the plane that maps $Z(f_3)$ onto itself, this correspondence is not one to one. If $\omega_1^{12} = \omega_2^{12} = 1$, then ω_1 and ω_2 will correspond to projectively equivalent monoids if and only if $\omega_1^3 = \omega_2^3$ or $\omega_1^3 \omega_2^3 = 1$. This means that there are three different quartic monoids with one $T_{3,3,4}$ singularity and one A_{11} singularity. One corresponds to those ω where $\omega^3 = 1$, one to those ω where $\omega^3 = -1$, and one to those ω where $\omega^6 = -1$. The first two of these have real representatives, $\omega = \pm 1$.

It is easy to see that for any set of multiplicities $m_1 + \dots + m_r = 12$, we can find real points $p_1, \dots, p_r$ such that condition (3) is satisfied. This completely classifies the possible configurations of singularities when f_3 is an irreducible nodal curve.

Case 2. The tangent cone is a cuspidal curve, and we can assume $f_3(x_1, x_2, x_3) = x_1^3 - x_2^2 x_3$. The cuspidal curve is singular at $(0 : 0 : 1)$ and can be parameterized by θ as a map from $\mathbb{P}^1$ to $\mathbb{P}^2$ where $\theta(s, t) = (s^2 t, s^3, t^3)$. The intersection numbers are determined by the degree 12 polynomial $f_4(\theta)$. As in the previous case, $f_4(\theta) \equiv 0$ if and only if f_3 is a factor of f_4 , and we will assume this is not the case. The multiplicity m of the factor s in $f_4(\theta)$ determines the type of singularity at O . If $m = 0$ (no factor s), then O is a Q_{10} singularity. If $m = 2$ or $m = 3$, then O is of type Q_{9+m} . If $m > 3$, then $(0 : 0 : 1)$ is a singular point on $Z(f_4)$, so the monoid has a singular line and is not considered in this article. Also, $m = 1$ is not possible, since $f_4(\theta(s, t)) = f_4(s^2 t, s^3, t^3)$ cannot contain st^{11} as a factor.

For each $m = 0, 2, 3$ we can analyze the possible configurations of other singularities on the monoid. Similarly to the previous case, any choice of parameter points $p_1, \dots, p_r \in \mathbb{P}^1 \setminus \{(0 : 1)\}$ and positive multiplicities $m_1, \dots, m_r$ with $\sum m_i = 12 - m$ corresponds, up to a nonzero constant, to a unique degree 12 polynomial q .

When $m = 2$ or $m = 3$, for any choice of parameter values and associated multiplicities, we can find a four dimensional family $f_4 = \lambda f'_4 + \ell f_3$ with the prescribed roots in $f_4(\theta)$. As before, the family gives projectively equivalent monoids.

When $m = 0$, one condition must be satisfied for q to be of the form $f_4(\theta)$, namely $b_{11} = 0$, where b_{11} is the coefficient of st^{11} in q .

For example, we can have an A_{11} singularity only if q is of the form $(\alpha s - \beta t)^{12}$. The condition $b_{11} = 0$ implies that either $q = \lambda s^{12}$ or $q = \lambda t^{12}$. The first case gives a surface with a singular line, while the other gives a monoid with an A_{11} singularity (see Figure 2). The line from O to the A_{11} singularity corresponds to the inflection point of $Z(f_3)$.

For any set of multiplicities $m_1, \dots, m_r$ with $m_1 + \dots + m_r = 12$, it is not hard to see that there exist real points $p_1, \dots, p_r$ such that the condition $b_{11} = 0$ is satisfied. It suffices to take $p_i = (\alpha_i : 1)$, with $\sum m_i \alpha_i = 0$ (the condition corresponding to $b_{11} = 0$). This completely classifies the possible configurations of singularities when f_3 is a cuspidal curve.

Case 3. The tangent cone is the product of a conic and a line that is not tangent to the conic, and we can assume $f_3 = x_3(x_1x_2 + x_3^2)$. Then $Z(f_3)$ is singular at $(1 : 0 : 0)$ and $(0 : 1 : 0)$, the intersections of the conic $Z(x_1x_2 + x_3^2)$ and the line $Z(x_3)$. For each f_4 we can associate four integers:

$$\begin{aligned} j_0 &:= I_{(1:0:0)}(x_1x_2 + x_3^2, f_4), \quad k_0 := I_{(1:0:0)}(x_3, f_4), \\ j_1 &:= I_{(0:1:0)}(x_1x_2 + x_3^2, f_4), \quad k_1 := I_{(0:1:0)}(x_3, f_4). \end{aligned}$$

We see that $k_0 > 0 \Leftrightarrow f_4(1 : 0 : 0) = 0 \Leftrightarrow j_0 > 0$, and that $Z(f_4)$ is singular at $(1 : 0 : 0)$ if and only if k_0 and j_0 both are bigger than one. These cases imply a singular line on the monoid, and are not considered in this article. The same holds for k_1, j_1 and the point $(0 : 1 : 0)$.

Define $r_i = \max(j_i, k_i)$ for $i = 1, 2$. Then, by [19], O will be a singularity of type $T_{3,4+r_0,4+r_1}$ if $r_0 \leq r_1$, or of type $T_{3,4+r_1,4+r_0}$ if $r_0 \geq r_1$.

We can parameterize the line $Z(x_3)$ by θ_1 where $\theta_1(s, t) = (s, t, 0)$, and the conic $Z(x_1x_2 + x_3^2)$ by θ_2 where $\theta_2(s, t) = (s^2, -t^2, st)$. Similarly to the previous cases, roots of $f_4(\theta_1)$ correspond to intersections between $Z(f_4)$ and the line $Z(x_3)$, while roots of $f_4(\theta_2)$ correspond to intersections between $Z(f_4)$ and the conic $Z(x_1x_2 + x_3^2)$.

For any legal values of j_0, j_1, k_0 and k_1 , parameter points

$$(\alpha_1 : \beta_1), \dots, (\alpha_{m_r} : \beta_{m_r}) \in \mathbb{P}^1 \setminus \{(0 : 1), (1 : 0)\},$$

with multiplicities $m_1, \dots, m_r$ such that $m_1 + \dots + m_r = 4 - k_0 - k_1$, and parameter points

$$(\alpha'_1 : \beta'_1), \dots, (\alpha'_{m'_r} : \beta'_{m'_r}) \in \mathbb{P}^1 \setminus \{(0 : 1), (1 : 0)\},$$

with multiplicities $m'_1, \dots, m'_{r'}$ such that $m'_1 + \dots + m'_{r'} = 8 - j_0 - j_1$, we can fix polynomials q_1 and q_2 such that

- q_1 is nonzero, of degree 4, and has factors s^{k_1}, t^{k_0} and $(\beta_i s - \alpha_i t)^{m_i}$ for $i = 1, \dots, r$,
- q_2 is nonzero, of degree 8, and has factors s^{j_1}, t^{j_0} and $(\beta'_i s - \alpha'_i t)^{m'_i}$ for $i = 1, \dots, r'$.

Now q_1 and q_2 are determined up to multiplication by nonzero constants. Write $q_1 = b_0 s^4 + \dots + b_4 t^4$ and $q_2 = c_0 s^8 + \dots + c_8 t^8$.

The classification of singularities on the monoid consists of describing the conditions on the parameter points and nonzero constants λ_1 and λ_2 for the pair $(\lambda_1 q_1, \lambda_2 q_2)$ to be on the form $(f_4(\theta_1), f_4(\theta_2))$ for some f_4 .

Similarly to the previous cases, $f_4(\theta_1) \equiv 0$ if and only if x_3 is a factor in f_4 and $f_4(\theta_2) \equiv 0$ if and only if $x_1x_2 + x_3^2$ is a factor in f_4 . Since $f_3 = x_3(x_1x_2 + x_3^2)$, both cases will make the monoid reducible, so we only consider $\lambda_1, \lambda_2 \neq 0$.

We use linear algebra to study the relationship between the coefficients $a_1 \dots a_{15}$ of f_4 and the polynomials q_1 and q_2 . We find $(\lambda_1 q_1, \lambda_2 q_2)$ to be of the form $(f_4(\theta_1), f_4(\theta_2))$ if and only if $\lambda_1 b_0 = \lambda_2 c_0$ and $\lambda_1 b_4 = \lambda_2 c_8$. Furthermore, the pair $(\lambda_1 q_1, \lambda_2 q_2)$ will fix f_4 modulo f_3 . Since f_4 and λf_4 correspond to projectively equivalent monoids for any $\lambda \neq 0$, it is the ratio λ_1/λ_2 , and not λ_1 and λ_2 , that is important.

Recall that $k_0 > 0 \Leftrightarrow j_0 > 0$ and $k_1 > 0 \Leftrightarrow j_1 > 0$. If $k_0 > 0$ and $k_1 > 0$, then $b_0 = c_0 = b_4 = c_8 = 0$, so for any $\lambda_1, \lambda_2 \neq 0$ we have $(\lambda_1 q_1, \lambda_2 q_2) = (f_4(\theta_1), f_4(\theta_2))$ for some f_4 . Varying λ_1/λ_2 will give a one-parameter family of monoids for each choice of multiplicities and parameter points.

If $k_0 = 0$ and $k_1 > 0$, then $b_0 = c_0 = 0$. The condition $\lambda_1 b_4 = \lambda_2 c_8$ implies $\lambda_1/\lambda_2 = c_8/b_4$. This means that any choice of multiplicities and parameter points will give a unique monoid up to projective equivalence. The same goes for the case where $k_0 > 0$ and $k_1 = 0$.

Finally, consider the case where $k_0 = k_1 = 0$. For $(\lambda_1 q_1, \lambda_2 q_2)$ to be of the form $(f_4(\theta_1), f_4(\theta_2))$ we must have $\lambda_1/\lambda_2 = c_8/b_4 = c_0/b_0$. This translates into a condition on the parameter points, namely

$$\frac{(\beta'_1)^{m'_1} \cdots (\beta'_{r'})^{m'_{r'}}}{\beta_1^{m_1} \cdots \beta_r^{m_r}} = \frac{(\alpha'_1)^{m'_1} \cdots (\alpha'_{r'})^{m'_{r'}}}{\alpha_1^{m_1} \cdots \alpha_r^{m_r}}. \quad (4)$$

In other words, if condition (4) holds, we have a unique monoid up to projective equivalence.

It is easy to see that for any choice of multiplicities, it is possible to find real parameter points such that condition (4) is satisfied. This completes the classification of possible singularities when the tangent cone is a conic plus a chordal line.

Case 4. The tangent cone is the product of a conic and a line tangent to the conic, and we can assume $f_3 = x_3(x_1 x_3 + x_2^2)$. Now $Z(f_3)$ is singular at $(1 : 0 : 0)$. For each f_4 we can associate two integers

$$j_0 := I_{(1:0:0)}(x_1 x_3 + x_2^2, f_4) \quad \text{and} \quad k_0 := I_{(1:0:0)}(x_3, f_4).$$

We have $j_0 > 0 \Leftrightarrow k_0 > 0$, $j_0 > 1 \Leftrightarrow k_0 > 1$. Furthermore, j_0 and k_0 are both greater than 2 if and only if $Z(f_4)$ is singular at $(1 : 0 : 0)$, a case we have excluded. The singularity at O will be of the S series, from [1], [2].

We can parameterize the conic $Z(x_1 x_3 + x_2^2)$ by θ_2 and the line $Z(x_3)$ by θ_1 where $\theta_2(s : t) = (s^2 : st : -t^2)$ and $\theta_1(s : t) = (s : t : 0)$. As in the previous case, the monoid is reducible if and only if $f_4(\theta_1) \equiv 0$ or $f_4(\theta_2) \equiv 0$. Consider two nonzero polynomials

$$\begin{aligned} q_1 &= b_0 s^4 + b_1 s^3 t + b_2 s^2 t^2 + b_3 s t^3 + b_4 t^4 \\ q_2 &= c_0 s^8 + c_1 s^7 t + \cdots + c_7 s t^7 + c_8 t^8. \end{aligned}$$

Now $(\lambda_1 q_1, \lambda_2 q_2) = (f_4(\theta_1), f_4(\theta_2))$ for some f_4 if and only if $\lambda_1 b_0 = \lambda_2 c_0$ and $\lambda_1 b_1 = \lambda_2 c_1$. As before, only the cases where $\lambda_1, \lambda_2 \neq 0$ are interesting. We see that $(\lambda_1 q_1, \lambda_2 q_2) = (f_4(\theta_1), f_4(\theta_2))$ for some $\lambda_1, \lambda_2 \neq 0$ if and only if the following hold:

- $b_0 = 0 \Leftrightarrow c_0 = 0$ and $b_1 = 0 \Leftrightarrow c_1 = 0$
- $b_0 c_1 = b_1 c_0$.

The classification of other singularities (than O) is very similar to the previous case. Roots of $f_4(\theta_1)$ and $f_4(\theta_2)$ away from $(1 : 0)$ correspond to intersections of $Z(f_3)$ and $Z(f_4)$ away from the singular point of $Z(f_3)$, and when one such intersection is multiple, there is a corresponding singularity on the monoid.

Now assume $(\lambda_1 q_1, \lambda_2 q_2) = (f_4(\theta_1), f_4(\theta_2))$ for some $\lambda_1, \lambda_2 \neq 0$ and some f_4 . If $b_0 \neq 0$ (equivalent to $c_0 \neq 0$) then $j_0 = k_0 = 0$ and $\lambda_1/\lambda_2 = c_0/b_0$. If $b_0 = c_0 = 0$ and $b_1 \neq 0$ (equivalent to $c_1 \neq 0$), then $j_0 = k_0 = 1$, and $\lambda_1/\lambda_2 = c_1/b_1$. If $b_0 = b_1 = c_0 = c_1 = 0$, then $j_0, k_0 > 1$ and any value of λ_1/λ_2 will give $(\lambda_1 q_1, \lambda_2 q_2)$ of the form $(f_4(\theta_1), f_4(\theta_2))$ for some f_4 . Thus we get a one-dimensional family of monoids for this choice of q_1 and q_2 .

Now consider the possible configurations of other singularities on the monoid. Assume that $j'_0 \leq 8$ and $k'_0 \leq 4$ are nonnegative integers such that $j_0 > 0 \Leftrightarrow k_0 > 0$ and $j_0 > 1 \Leftrightarrow k_0 > 1$. For any set of multiplicities $m_1, \dots, m_r$ with $m_1 + \dots + m_r = 4 - k'_0$ and $m'_1, \dots, m'_r$ with $m'_1 + \dots + m'_r = 8 - j'_0$, there exists a polynomial f_4 with real coefficients such that $f_4(\theta_1)$ has real roots away from $(1 : 0)$ with multiplicities $m_1, \dots, m_r$, and $f_4(\theta_2)$ has real roots away from $(1 : 0)$ with multiplicities $m'_1, \dots, m'_r$. Furthermore, for this f_4 we have $k_0 = k'_0$ and $j_0 = j'_0$. Proposition 6 will give the singularities that occur in addition to O .

This completes the classification of the singularities on a quartic monoid (other than O) when the tangent cone is a conic plus a tangent.

Case 5. The tangent cone is three general lines, and we assume $f_3 = x_1 x_2 x_3$.

For each f_4 we associate six integers,

$$\begin{aligned} k_2 &:= I_{(1:0:0)}(f_4, x_2), \quad l_1 := I_{(0:1:0)}(f_4, x_1), \quad m_1 := I_{(0:0:1)}(f_4, x_1), \\ k_3 &:= I_{(1:0:0)}(f_4, x_3), \quad l_3 := I_{(0:1:0)}(f_4, x_3), \quad m_2 := I_{(0:0:1)}(f_4, x_2). \end{aligned}$$

Now $k_2 > 0 \Leftrightarrow k_3 > 0$, $l_1 > 0 \Leftrightarrow l_3 > 0$, and $m_1 > 0 \Leftrightarrow m_2 > 0$. If both k_2 and k_3 are greater than 1, then the monoid has a singular line, a case we have excluded. The same goes for the pairs (l_1, l_3) and (m_1, m_2) .

When the monoid does not have a singular line, we define $j_k = \max(k_2, k_3)$, $j_l = \max(l_1, l_3)$ and $j_m = \max(m_1, m_2)$. If $j_k \leq j_l \leq j_m$, then [19] gives that O is a $T_{4+j_k, 4+j_l, 4+j_m}$ singularity.

The three lines $Z(x_1)$, $Z(x_2)$ and $Z(x_3)$ are parameterized by θ_1 , θ_2 and θ_3 where $\theta_1(s : t) = (0 : s : t)$, $\theta_2(s : t) = (s : 0 : t)$ and $\theta_3(s : t) = (s : t : 0)$. Roots of the polynomial $f_4(\theta_i)$ away from $(1 : 0)$ and $(0 : 1)$ correspond to intersections between $Z(f_4)$ and $Z(x_i)$ away from the singular points of $Z(f_3)$.

As before, we are only interested in the cases where none of $f_4(\theta_i) \equiv 0$ for $i = 1, 2, 3$, as this would make the monoid reducible.

For the study of other singularities on the monoid we consider nonzero polynomials

$$\begin{aligned} q_1 &= b_0 s^4 + b_1 s^3 t + b_2 s^2 t^2 + b_3 s t^3 + b_4 t^4, \\ q_2 &= c_0 s^4 + c_1 s^3 t + c_2 s^2 t^2 + c_3 s t^3 + c_4 t^4, \\ q_3 &= d_0 s^4 + d_1 s^3 t + d_2 s^2 t^2 + d_3 s t^3 + d_4 t^4. \end{aligned}$$

Linear algebra shows that $(\lambda_1 q_1, \lambda_2 q_2, \lambda_3 q_3) = (f_4(\theta_1), f_4(\theta_2), f_4(\theta_3))$ for some f_4 if and only if $\lambda_1 b_0 = \lambda_3 d_4$, $\lambda_1 b_4 = \lambda_2 c_4$, and $\lambda_2 c_0 = \lambda_3 d_0$. A simple analysis shows the following: There exist $\lambda_1, \lambda_2, \lambda_3 \neq 0$ such that

$$(\lambda_1 q_1, \lambda_2 q_2, \lambda_3 q_3) = (f_4(\theta_1), f_4(\theta_2), f_4(\theta_3))$$

for some f_4 , and such that $Z(f_4)$ and $Z(f_3)$ have no common singular point if and only if all of the following hold:

- $b_0 = 0 \leftrightarrow d_4 = 0$ and $b_0 = d_4 = 0 \rightarrow (b_1 \neq 0 \text{ or } d_3 \neq 0)$,
- $b_4 = 0 \leftrightarrow c_4 = 0$ and $b_4 = c_4 = 0 \rightarrow (b_3 \neq 0 \text{ or } c_3 \neq 0)$,
- $c_0 = 4 \leftrightarrow d_0 = 0$ and $c_0 = d_0 = 0 \rightarrow (c_1 \neq 0 \text{ or } d_1 \neq 0)$,
- $b_0 c_4 d_0 = b_4 c_0 d_4$.

Similarly to the previous cases we can classify the possible configurations of other singularities by varying the multiplicities of the roots of the polynomials q_1 , q_2 and q_3 . Only the multiplicities of the roots $(0 : 1)$ and $(1 : 0)$ affect the first three bullet points above. Then, for any set of multiplicities of the rest of the roots, we can find q_1 , q_2 and q_3 such that the last bullet point is satisfied. This completes the classification when $Z(f_3)$ is the product of three general lines.

Case 6. The tangent cone is three lines meeting in a point, and we can assume that $f_3 = x_2^3 - x_2 x_3^2$. We write $f_3 = \ell_1 \ell_2 \ell_3$ where $\ell_1 = x_2$, $\ell_2 = x_2 - x_3$ and $\ell_3 = x_2 + x_3$, representing the three lines going through the singular point $(1 : 0 : 0)$. For each f_4 we associate three integers j_1 , j_2 and j_3 defined as the intersection numbers $j_i = I_{(1:0:0)}(f_4, \ell_i)$. We see that $j_1 = 0 \Leftrightarrow j_2 = 0 \Leftrightarrow j_3 = 0$, and that $Z(f_4)$ is singular at $(1 : 0 : 0)$ if and only if two of the integers j_1 , j_2 , j_3 are greater than one. (Then all of them will be greater than one.) The singularity will be of the U series [1], [2].

The three lines $Z(\ell_1)$, $Z(\ell_2)$ and $Z(\ell_3)$ can be parameterized by θ_1 , θ_2 , and θ_3 where $\theta_1(s : t) = (s : 0 : t)$, $\theta_2(s : t) = (s : t : t)$ and $\theta_3(s : t) = (s : t : -t)$.

For the study of other singularities on the monoid we consider nonzero polynomials

$$\begin{aligned} q_1 &= b_0 s^4 + b_1 s^3 t + b_2 s^2 t^2 + b_3 s t^3 + b_4 t^4, \\ q_2 &= c_0 s^4 + c_1 s^3 t + c_2 s^2 t^2 + c_3 s t^3 + c_4 t^4, \\ q_3 &= d_0 s^4 + d_1 s^3 t + d_2 s^2 t^2 + d_3 s t^3 + d_4 t^4. \end{aligned}$$

Linear algebra shows that $(\lambda_1 q_1, \lambda_2 q_2, \lambda_3 q_3) = (f_4(\theta_1), f_4(\theta_2), f_4(\theta_3))$ for some f_4 if and only if $\lambda_1 b_0 = \lambda_2 c_4 = \lambda_3 d_0$, and $2\lambda_1 b_1 = \lambda_2 c_1 + \lambda_3 d_1$. There exist $\lambda_1, \lambda_2, \lambda_3 \neq 0$ such that $(\lambda_1 q_1, \lambda_2 q_2, \lambda_3 q_3) = (f_4(\theta_1), f_4(\theta_2), f_4(\theta_3))$ for some f_4 and such that $Z(f_4)$ and $Z(f_3)$ have no common singular point if and only if all of the following hold:

- $b_0 = 0 \leftrightarrow c_0 = 0 \leftrightarrow d_0 = 0$,
- if $b_0 = c_0 = d_0 = 0$, then at least two of b_1 , c_1 , and d_1 are different from zero,
- $2b_1 c_0 d_0 = b_0 c_1 d_0 + b_0 c_0 d_1$.

As in all the previous cases we can classify the possible configurations of other singularities for all possible j_1, j_2, j_3 . As before, the first bullet point only affect the multiplicity of the factor t in q_1 , q_2 and q_3 . For any set of multiplicities for the rest of the roots, we can find q_1 , q_2 , q_3 with real roots of the given multiplicities such that the last bullet point is satisfied. This completes the classification of the singularities (other than O) when $Z(f_3)$ is three lines meeting in a point.

Case 7. The tangent cone is a double line plus a line, and we can assume $f_3 = x_2 x_3^2$. The tangent cone is singular along the line $Z(x_3)$. The line $Z(x_2)$ is parameterized by θ_1 and the line $Z(x_3)$ is parameterized by θ_2 where $\theta_1(s : t) = (s : 0 : t)$ and $\theta_2(s : t) = (s : t : 0)$. The monoid is reducible if and only if $f_4(\theta_1)$ or $f_4(\theta_2)$ is identically zero, so we assume that neither is identically zero. For each f_4 we associate

two integers, $j_0 := I_{(1:0:0)}(f_4, x_2)$ and $k_0 := I_{(1:0:0)}(f_4, x_3)$. Furthermore, we write $f_4(\theta_2)$ as a product of linear factors

$$f_4(\theta_2) = \lambda s^{k_0} \prod_{i=0}^r (\alpha_i s - t)^{m_i}.$$

Now the singularity at O will be of the V series and depends on j_0 , k_0 and $m_1, \dots, m_r$.

Other singularities on the monoid correspond to intersections of $Z(f_4)$ and the line $Z(x_2)$ away from $(1 : 0 : 0)$. Each such intersection corresponds to a root in the polynomial $f_4(\theta_1)$ different from $(1 : 0)$. Let $j'_0 \leq 4$ and $k'_0 \leq 4$ be integers such that $j_0 > 0 \leftrightarrow k_0 > 0$. Then, for any homogeneous polynomials q_1, q_2 in s, t of degree 4 such that s is a factor of multiplicity j'_0 in q_1 and of multiplicity k'_0 in q_2 , there is a polynomial f_4 and nonzero constants λ_1 and λ_2 such that $k_0 = k'_0$, $j_0 = j'_0$ and $(\lambda_1 q_1, \lambda_2 q_2) = (f_4(\theta_1), f_4(\theta_2))$. Furthermore, if q_1 and q_2 have real coefficients, then f_4 can be selected with real coefficients. This follows from an analysis similar to case 5 and completes the classification of singularities when the tangent cone is a product of a line and a double line.

Case 8. The tangent cone is a triple line, and we assume that $f_3 = x_3^3$. The line $Z(x_3)$ is parameterized by θ where $\theta(s, t) = (s, t, 0)$. Assume that the polynomial $f_4(\theta)$ has r distinct roots with multiplicities $m_1, \dots, m_r$. (As before $f_4(\theta) \equiv 0$ if and only if the monoid is reducible.) Then the type of the singularity at O will be of the V' series [3, p. 267]. The integers $m_1, \dots, m_r$ are constant under right equivalence over $\mathbb{C}$. Note that one can construct examples of monoids that are right equivalent over $\mathbb{C}$, but not over $\mathbb{R}$ (see Figure 4).




Fig. 4. The monoids $Z(z^3 + xy^3 + x^3y)$ and $Z(z^3 + xy^3 - x^3y)$ are right equivalent over $\mathbb{C}$ but not over $\mathbb{R}$.

The tangent cone is singular everywhere, so there can be no other singularities on the monoid.

Case 9. The tangent cone is a smooth cubic curve, and we write $f_3 = x_1^3 + x_2^3 + x_3^3 + 3ax_1x_2x_3$ where $a^3 \neq -1$. This is a one-parameter family of elliptic curves, so we cannot use the parameterization technique of the other cases. The singularity at O will be a P_8 singularity (cf. [3, p. 185]), and other singularities correspond to intersections between $Z(f_3)$ and $Z(f_4)$, as described by Proposition 6.

To classify the possible configurations of singularities on a monoid with a nonsingular (projective) tangent cone, we need to answer the following question: For any positive integers $m_1, \dots, m_r$ such that $\sum_{i=1}^r m_i = 12$, does there, for *some* $a \in \mathbb{R} \setminus \{-1\}$, exist a polynomial f_4 with real coefficients such that $Z(f_3, f_4) = \{p_1, \dots, p_r\} \in \mathbb{P}^2(\mathbb{R})$ and $I_{p_i}(f_3, f_4) = m_i$ for $i = 1, \dots, r$? Rohn [15, p. 63] says that one can always find curves $Z(f_3), Z(f_4)$ with this property. Here we shall show that for *any* $a \in \mathbb{R} \setminus \{-1\}$ we can find a suitable f_4 .

In fact, in almost all cases f_4 can be constructed as a product of linear and quadratic terms in a simple way. The difficult cases are $(m_1, m_2) = (11, 1)$, $(m_1, m_2, m_3) = (8, 3, 1)$, and $(m_1, m_2) = (5, 7)$. For example, the case where $(m_1, m_2, m_3) = (3, 4, 5)$ can be constructed as follows: Let $f_4 = \ell_1 \ell_2 \ell_3^2$ where ℓ_1 and ℓ_2 define tangent lines at inflection points p_1 and p_3 of $Z(f_3)$. Let ℓ_3 define a line that intersects $Z(f_3)$ once at p_3 and twice at another point p_2 . Note that the points p_1, p_2 and p_3 can be found for any $a \in \mathbb{R} \setminus \{-1\}$.

The case $(m_1, m_2) = (11, 1)$ is also possible for every $a \in \mathbb{R} \setminus \{-1\}$. For any point p on $Z(f_3)$ there exists an f_4 such that $I_p(f_3, f_4) \geq 11$. For all except a finite number of points, we have equality [11], so the case $(m_1, m_2) = (11, 1)$ is possible for any $a \in \mathbb{R} \setminus \{-1\}$. The case $(m_1, m_2, m_3) = (8, 3, 1)$ is similar, but we need to let f_4 be a product of the tangent at an inflection point with another cubic.

The case $(m_1, m_2) = (5, 7)$ is harder. Let $a = 0$. Then we can construct a conic C that intersects $Z(f_3)$ with multiplicity five in one point and multiplicity one in an inflection point, and choosing $Z(f_4)$ as the union of C and twice the tangent line through the inflection point will give the desired example. The same can be done for $a = -4/3$. By using the computer algebra system SINGULAR [6] we can show that these constructions can be continuously extended to any $a \in \mathbb{R} \setminus \{-1\}$. This completes the classification of singularities on a monoid when the tangent cone is smooth.

In the Cases 3, 5, and 6, not all real equations of a given type can be transformed to the chosen forms by a real transformation.

In Case 3 the conic may not intersect the line in two real points, but rather in two complex conjugate points. Then we can assume $f_3 = x_3(x_1x_3 + x_1^2 + x_2^2)$, and the singular points are $(1 : \pm i : 0)$. For any real f_4 , we must have

$$I_{(1:i:0)}(x_1x_3 + x_1^2 + x_2^2, f_4) = I_{(1:-i:0)}(x_1x_3 + x_1^2 + x_2^2, f_4)$$

and

$$I_{(1:i:0)}(x_3, f_4) = I_{(1:-i:0)}(x_3, f_4),$$

so only the cases where $j_0 = j_1$ and $k_0 = k_1$ are possible. Apart from that, no other restrictions apply.

In Case 5, two of the lines can be complex conjugate, and we assume $f_3 = x_3(x_1^2 + x_2^2)$. A configuration from the previous analysis is possible for real coefficients of f_4 if

and only if $m_1 = m_2$, $k_2 = l_1$, and $k_3 = l_3$. Furthermore, only the singularities that correspond to the line $Z(x_3)$ will be real.

In Case 6, two of the lines can be complex conjugate, and then we may assume $f_3 = x_2^3 + x_3^3$. Now, if j_3 denotes the intersection number of $Z(f_4)$ with the real line $Z(x_2 + x_3)$, precisely the cases where $j_1 = j_2$ are possible, and only intersections with the line $Z(x_2 + x_3)$ may contribute to real singularities.

This concludes the classification of real and complex singularities on real monoids of degree 4.

Remark. In order to describe the various monoid singularities, Rohn [15] computes the “class reduction” due to the presence of the singularity, in (almost) all cases. (The class is the degree of the dual surface [14, p. 262].) The class reduction is equal to the local intersection multiplicity of the surface with two general polar surfaces. This intersection multiplicity is equal to the sum of the Milnor number and the Milnor number of a general plane section through the singular point [20, Cor. 1.5, p. 320]. It is not hard to see that a general plane section has either a D_4 (Cases 1–6, 9), D_5 (Case 7), or E_6 (Case 8) singularity. Therefore one can retrieve the Milnor number of each monoid singularity from Rohn’s work.

Acknowledgements

We would like to thank the referees for helpful comments. This research was supported by the European Union through the project IST 2001–35512 ‘Intersection algorithms for geometry based IT applications using approximate algebraic methods’ (GAIA II).

References

1. V. I. Arnol’d. Normal forms of functions in neighbourhoods of degenerate critical points. *Russian Math Surveys*, 29ii:10–50, 1974.
2. V. I. Arnol’d. Critical points of smooth functions and their normal forms. *Russian Math Surveys*, 30v:1–75, 1975.
3. V. I. Arnol’d, S. M. Gusein-Zade, and A. N. Varchenko. *Singularities of differentiable maps. Vol. I*, volume 82 of *Monographs in Mathematics*. The classification of critical points, caustics and wave fronts, Translated from the Russian by Ian Porteous and Mark Reynolds. Birkhäuser Boston Inc., Boston, MA, 1985.
4. J. W. Bruce and C. T. C. Wall. On the classification of cubic surfaces. *J. London Math. Soc.* (2), 19(2):245–256, 1979.
5. S. Endraß et al. SURF 1.0.4. 2003. A Computer Software for Visualising Real Algebraic Geometry, <http://surf.sourceforge.net>.
6. G.-M. Greuel, G. Pfister, and H. Schönemann. SINGULAR 3.01. A Computer Algebra System for Polynomial Computations. Centre for Computer Algebra, University of Kaiserslautern (2005). <http://www.singular.uni-kl.de>.
7. C. M. Jessop. Quartic surfaces with singular points. *Cambridge University Press*, 1916.
8. Horst Knörrer and Thomas Miller. Topologische Typen reeller kubischer Flächen. *Math. Z.*, 195(1):51–67, 1987.
9. Oliver Labs and Duco van Straten. The Cubic Surface Homepage. 2001. <http://www.Cubics.AlgebraicSurface.net>.

10. Henry B. Laufer. On minimally elliptic singularities. *Amer. J. Math.*, 99(6):1257–1295, 1977.
11. Claire Moura. Local intersections of plane algebraic curves. *Proc. Amer. Math. Soc.*, 132(3):687–690 (electronic), 2004.
12. Sonia Pérez-Díaz, Juana Sendra, and J. Rafael Sendra. Parametrization of approximate algebraic curves by lines. *Theoret. Comput. Sci.*, 315(2-3):627–650, 2004.
13. Sonia Pérez-Díaz, Juana Sendra, and J. Rafael Sendra. Parametrization of approximate algebraic surfaces by lines. *Comput. Aided Geom. Design*, 22(2):147–181, 2005.
14. Ragni Piene. Polar classes of singular varieties. *Ann. Sci. École Norm. Sup. (4)*, 11(2):247–276, 1978.
15. Karl Rohn. Ueber die Flächen vierter Ordnung mit dreifachem Punkte. *Math. Ann.*, 24(1):55–151, 1884.
16. L. Schläfli. On the distribution of surfaces of the third order into species. *Philos. Trans. Royal Soc.*, CLIII:193–241, 1863.
17. Thomas W. Sederberg, Jianmin Zheng, Kris Klimaszewski, and Tor Dokken. Approximate implicitization using monoid curves and surfaces. *Graphical Models and Image Processing*, 61(4):177–198, 1999.
18. B. Segre. *The Non-singular Cubic Surfaces*. Oxford University Press, Oxford, 1942.
19. Tadashi Takahashi, Kimio Watanabe, and Teiichi Higuchi. On the classification of quartic surfaces with triple point. I, II. *Sci. Rep. Yokohama Nat. Univ. Sect. I*, (29):47–70, 71–94, 1982.
20. Bernard Teissier. Cycles évanescents, sections planes et conditions de Whitney. In *Singularités à Cargèse (Rencontre Singularités Géom. Anal., Inst. Études Sci., Cargèse, 1972)*, pages 285–362. Astérisque, Nos. 7 et 8. Soc. Math. France, Paris, 1973.
21. Yumiko Umezū. On normal projective surfaces with trivial dualizing sheaf. *Tokyo J. Math.*, 4(2):343–354, 1981.